



Soluções dos exercícios de Primitivas e Cálculo Integral

Exercício 1. Primitivas:

$$(a) x^4 + \frac{x^3}{3} - \frac{x^2}{2} + x + c;$$

$$(c) -\frac{1}{x} - 2\sqrt{x} - \frac{1}{15} \sin(5x) + c;$$

$$(e) -\ln|\cos(x)| + \ln|1+x| + c;$$

$$(g) -e^{\cos(x)} + c;$$

$$(i) \frac{1}{12}(x^3+1)^4 + c;$$

$$(k) \frac{\arctan^2 x}{2} + \ln|\arctan x| + c;$$

$$(m) \ln|1+\sin(x)| + \arctan(\sin(x)) + c;$$

$$(o) 2x - 2\ln|(x+3)(x+7)| + c;$$

$$(q) x + \frac{5}{2} \ln(x^2+1) - 3\arctan(x) + c;$$

$$(s) -\frac{1}{5}\sqrt{2-5x^2} - \frac{1}{\sqrt{5}} \arccos\left(\sqrt{\frac{5}{2}}x\right) + c;$$

$$(b) -\frac{\cos(3x)}{3} - \frac{1}{12}x^6 + \frac{2}{3}\sqrt[3]{x^3} + c;$$

$$(d) \frac{3}{2}\sqrt[3]{x^2} - \frac{1}{3}e^{3x} - 2\arctan(x) - \frac{1}{5}e^{-2x} + c;$$

$$(f) -\frac{1}{24}(2-4x)^6 + c;$$

$$(h) \frac{1}{3}e^{x^3} - \frac{1}{2}\sin(x^2) + c;$$

$$(j) \frac{(1+\ln(x))^6}{6} + c;$$

$$(l) \ln(1+e^x) + \arctan(e^x) + c;$$

$$(n) \frac{1}{4}\sin^4(x) - \frac{\cos^2(x)}{2} + c;$$

$$(p) \frac{1}{2}\arctan(2x) + \frac{\sqrt{3}}{6}\arctan\left(\frac{2}{\sqrt{3}}x\right) + c;$$

$$(r) x - \frac{7}{2}\ln(x^2+9) - \frac{4}{3}\arctan\left(\frac{x}{3}\right) + c;$$

$$(t) -\frac{e^{-x^5}}{5} - e^{\frac{1}{x}} + c.$$

Exercício 2.

b.

$$\int \cos^2(x) dx = \frac{x}{2} + \frac{\sin(2x)}{4} + c$$

$$\int \sin^2(x) dx = \frac{x}{2} - \frac{\sin(2x)}{4} + c$$

c.

$$\int \cos^4(x) dx = \frac{3}{8}x + \frac{\sin(4x)}{32} + \frac{\sin(2x)}{4} + c$$

$$\int \sin^6(x) dx = -2^{-5} \left(\frac{\sin(6x)}{6} - \frac{3 \sin(4x)}{2} + \frac{15 \sin(2x)}{2} + 10x \right) + c$$

Exercício 3. Primitivas:

$$(a) \frac{x e^{2x}}{2} - \frac{e^{2x}}{4} + c;$$

$$(b) -\frac{x^2 \cos(2x)}{2} + \frac{x \sin(2x)}{2} + \frac{\cos(2x)}{4} + c;$$

$$(c) \frac{e^{3x}}{3} \left(x^2 + \frac{x}{3} + \frac{8}{9} \right) + c;$$

$$(d) x \arctan(3x) - \frac{\ln(1+9x^2)}{6} + c;$$

$$(e) x \ln(x) - x + c;$$

$$(f) x \arccos(x) - \sqrt{1-x^2} + c;$$

$$(g) x(\arccos(x) + \arcsin(x)) + c;$$

$$(h) x(\arctan(x) + \arctan(1/x)) + c;$$

$$(i) \frac{e^x}{5} \left(\sin(2x) - 2 \cos(2x) \right) + c;$$

$$(j) \frac{e^{2x}}{5} \left(\sin(x) + 2 \cos(x) \right) + c;$$

$$(k) \frac{2}{3} \sqrt{x^3} \ln(x) - \frac{4}{9} \sqrt{x^3} + c;$$

$$(l) x \ln^2(x) - 2x \ln(x) + 2x + c;$$

$$(m) \frac{x^3}{3} \arctan x - \frac{x^2}{6} + \frac{\ln(1+x^2)}{6} + c;$$

$$(n) \frac{x^2 e^{x^2}}{2} - \frac{e^{x^2}}{2} + c.$$

Exercício 4. Primitivas:

$$(a) 2 \ln|x-2| - \ln|x-1| + c;$$

$$(b) \frac{\ln|x-1|}{6} - \frac{\ln|x+1|}{2} + \frac{\ln|x+2|}{3} + c;$$

$$(c) \frac{\ln(x^2+4)}{2} + c;$$

$$(d) \frac{3}{2} \ln|x+1| - \frac{1}{4} \ln(x^2+1) + \frac{1}{2} \arctan(x) + c;$$

$$(e) \frac{x^2}{2} + 2 \ln|x-1| - \ln|x+1| + c;$$

$$(f) \frac{x^2}{2} + 2x + \ln(x^2+1) + 2 \arctan(x) + c.$$

Exercício 5. Primitivas :

$$(a) \frac{1}{2} \left(\arctan(x) + \frac{x}{1+x^2} \right) + c;$$

$$(b) \ln |1 + \cos(x)| - \ln |\cos(x)| + c;$$

$$(c) -\frac{\ln |1 - \sin(x)|}{4} + \frac{1}{2(1 - \sin(x))} + \frac{\ln |1 + \sin(x)|}{4} + c;$$

$$(d) 2 \frac{\sqrt{(e^x + 1)^3}}{3} - 2\sqrt{e^x + 1} + c;$$

$$(e) -2 \arccos \left(\frac{x}{2} \right) + \frac{x}{2} \sqrt{4 - x^2} + c;$$

$$(f) \frac{2}{3} t^{\frac{3}{2}} - \frac{3}{4} t^{\frac{4}{3}} + \frac{6}{7} t^{\frac{7}{6}} - t + \frac{6}{5} t^{\frac{5}{6}} - \frac{3}{2} t^{\frac{2}{3}} + 2t^{\frac{1}{2}} - 3t^{\frac{1}{3}} + 6t^{\frac{1}{6}} - \ln(t^{\frac{1}{6}} + 1) + c.$$

Exercício 6. b.

$$\int \sqrt{4+t^2} dt = \frac{(t + \sqrt{t^2 + 4})^2}{8} - \frac{2}{(t + \sqrt{t^2 + 4})^2} + 2 \ln \left(\frac{t + \sqrt{t^2 + 4}}{2} \right) + c$$

Exercício 7. Integrais:

$$(a) -3;$$

$$(b) 4;$$

$$(c) 0;$$

$$(d) 0;$$

$$(e) -1;$$

$$(f) \frac{b^2 - a^2}{2}.$$

Exercício 8. Áreas:

$$(a) \frac{36}{\sqrt{2}};$$

$$(b) \frac{128}{15} + 2\pi;$$

$$(c) \frac{1}{12};$$

$$(d) \frac{7}{48}.$$

Exercício 9. -

Exercício 10. $\lim_{x \rightarrow 0} M_f(x) = f(0).$

Exercício 11. -

Exercício 12. $I = \frac{\ln(3)}{6} - \frac{\ln(2)}{3} + \frac{\arctan(\sqrt{2})}{3\sqrt{2}}$.

Exercício 13. Soluções:

(a) $-\frac{1}{2}$;

(b) 4;

(c) $\frac{\pi^2}{8}$;

(d) -1;

(e) Divergente;

(f) $\frac{\pi}{2}$.

Um último desafio

Exercício 14. Respostas:

$$I_0 = \frac{\pi}{2}$$

$$I_1 = 1.$$

$$\int_0^{\frac{\pi}{2}} \sin^n(x) dx = \frac{(2k+1)!}{k \prod_{i=0}^{k-1} (2i+1)^2}, n = 2k+1, k \in \mathbb{N}_0.$$